

**EFFICACY OF HEXAFLUORINE(R) FOR EMERGENT  
DECONTAMINATION OF HYDROFLUORIC ACID EYE AND SKIN  
SPLASHES**

L. Mathieu\*, J. Nehles\*\*, J. Blomet\*, A.H. Hall+

\* Laboratoire PREVOR, Valmondois, France

\*\* Mannesmann Hoesch Prazisrohr, GmbH, Remscheid, Germany

+ Texas Tech University Health Sciences Center - El Paso, Department of  
Emergency Medicine, Division of Toxicology; Toxicology Consulting and Medical  
Translating Services, Inc., El Paso, TX, USA

Presented at the Semiconductor Safety Association Annual Meeting, Arlington,  
VA, 4/25-28/2000

Funding for translation of this article from the French and editing for publication in  
English was provided by Laboratoire PREVOR, Moulin de Verville F95760,  
Valmondois, France.

**Address for Correspondence:**

Alan H. Hall, M.D.

TCMTS, Inc.

3456 Oxcart Run Street

El Paso, TX 79936

USA

Telephone: (915) 856-9396

FAX: (915) 856-9956

Email: [ahalltoxic@netscape.net](mailto:ahalltoxic@netscape.net)

**ABSTRACT:** Hexafluorine(R) is an amphoteric, hypertonic, polyvalent compound for decontaminating hydrofluoric acid (HF) eye and skin splashes. In a German metallurgy facility during the period of 1994-1998, all eye or skin splashes with 40% HF alone or with a 6% HF/15% HNO(3) mixture were initially decontaminated with Hexafluorine(R) within the first 2 minutes following the splash at the accident site by the victims themselves or co-workers who witnessed the accident. Eleven workers using 40% HF or a 6% HF/15% HNO(3) mixture sustained eye ( 2 cases) or skin (10 cases) splashes (1 combined) during 1994-1998. Hexafluorine (R) was used within the first 2 minutes, and a second Hexafluorine(R) decontamination was done on arrival at the plant infirmary. No further medical or surgical treatment was needed, no workers developed chemical burns, and none had lost work time. Amphoteric Hexafluorine (R) binds both hydrogen and fluoride ions and has an affinity for fluoride 100 times greater than that of calcium gluconate. In experimental animals, Hexafluorine (R) prevented 70% HF skin burns, while water decontamination alone or followed by 2.5% calcium gluconate gel inunction did not. These 11 cases demonstrate the efficacy of Hexafluorine (R) in decontaminating HF or combined HF/HNO(3) splashes.

**Key Words:** Hydrofluoric acid; Hydrofluoric acid, burns; Hydrogen fluoride; Hydrogen fluoride, burns; Skin decontamination; Eye decontamination; Hexafluorine(R)

Hydrofluoric acid (HF) is a corrosive and toxic chemical and splashes can cause physical and psychological sequelae (1-3) and sometimes fatalities (4-9). The corrosive lesions are caused by penetration of  $H^+$  and  $F^-$  ions into the tissues; chelation of calcium leading to toxic systemic hypocalcemia (10-11) and other electrolyte imbalances (hypomagnesemia, hyperkalemia) may occur (12-13). Because of the risks presented by hydrofluoric acid and its widespread usage in industry (6), numerous publications have described a variety of treatment modalities for HF burns in both experimental animals (14-23) and in human accidental exposures (4,13,24).

Hexafluorine (R) is a novel amphoteric, hypertonic, polyvalent compound specifically developed for decontamination of hydrofluoric acid eye and skin splashes. The cases reported here demonstrate the ability of Hexafluorine (R) to decontaminate HF eye/skin splashes and to prevent the expected development of HF burns and their sequelae.

### **METHODS**

Mannesmann is a German metallurgical facility in which 40% HF is diluted and then mixed with nitric acid ( $HNO_3$ ) to produce a mixture of 6% HF/15%  $HNO_3$  utilized for treatment of stainless steel tubing used in the automobile industry for manufacturing luxury car exhaust pipes. Eye or skin splashes with 40% HF, or during maintenance operations.

In 1994, after experience with attempting decontamination of eye or skin splashes with 6% HF/15%  $HNO_3$  with water lavage followed by topical application of 2.5% calcium gluconate gel (skin splashes only) without complete success, the Mannesmann Medical and Health and Safety Services decided to train workers about the hazards of these chemicals and the use of Hexafluorine(R) decontamination and the various application devices which had been previously situated throughout the workplace.

During the period of 1994-1998, all eye or skin splashes with 40% HF alone or the 6% HF/15%  $HNO_3$  mixture were initially decontaminated with Hexafluorine(R) within the first 2 minutes

(between 30 and 120 seconds following the splash) at the accident site by the victims themselves or co-workers who witnessed the accident. Workers with skin splashes were undressed prior to decontamination.

The amount of Hexafluorine(R) used depended on the duration of contact and the extent of the total body surface area involved. For initial decontamination of eye splashes, 500 ml of Hexafluorine(R) was applied within the first 2 minutes (30-120 seconds) following exposure. For skin splashes, an automatic portable 5 L Hexafluorine (R) shower was used to decontaminate each exposed worker's entire body and utilized within 2 minutes (30-120 seconds) following exposure. In each instance, a second Hexafluorine (R) decontamination was carried out when the exposed worker arrived at the plant infirmary, after which clean work clothing was provided.

## **RESULTS**

During the period of 1994-1998, 11 workers sustained eye or skin splashes, 6 of which involved the 6% HF/15% HNO(3) mixture and 5 of which were with 40% HF. Exposed workers were all males aged 35 +/- 11 years. One worker sustained both eye and skin splashes with 40% HF and 1 worker had only an eye splash with the 6% HF/15 % HNO(3) mixture (Table 1).

The 10 skin splashes involved from 0.2% (1 finger) to 16.5% of the total body surface area (Table 2). In 6 cases, the total body surface area involved was equal to or greater than 4%. In one case each, splashes with either the 6% HF/15% HNO(3) mixture or 40% HF involved greater than 10% of the total body surface area, 10.5% and 16.5% respectively. Exposed areas included the hands, upper or lower extremities, face, eyes, and thorax.

No chemical burns or sequelae of any sort were observed in these 11 exposed workers. There was no requirement for any type of medical or surgical treatment other than Hexafluorine(R) decontamination and none of these workers had any lost work time.

## DISCUSSION

Numerous methods for decontamination and treating HF dermal burns can be found in the published literature (25,12,26-29). Splashes with concentrated HF (40-70%, anhydrous) rapidly produce very painful lesions (30), requiring that decontamination and treatment be undertaken immediately following exposure. In spite of early decontamination with water followed by repeated topical inunction or subcutaneous injection of calcium gluconate, development of burns often cannot be prevented. The risk of developing systemic HF toxicity (hypocalcemia, hypomagnesemia, hyperkalemia and their attendant risks) is greatest with concentrated HF splashes involving 2.5% or more of the total body surface area, which can sometimes be fatal (4). Splashes with dilute HF have been successfully decontaminated with water followed by topical application of calcium gluconate gel. In such cases, the difficulty lies in perceiving the need to immediately undertake these measures in the absence of pain, which may be delayed in onset for as much as 24-48 hours after exposure (30-32). The duration of the HF contact with the tissues may be prolonged in such situations; although the fluoride ion is initially bound to the calcium ion derived from calcium gluconate, it may later be released and can then initiate a burn. With dermal exposure to either dilute or concentrate HF in some cases surgical debridement, excision, or even amputation of necrotic areas may be required (33-36).

The relatively short duration of this study (1994-1998) plus the utilization of safety goggles resulted in inclusion of only 2 cases of HF eye splashes. In both cases, rapid decontamination with Hexafluorine (R) prevented ocular burns in 1 case each of eye splashes with concentrated or dilute HF.

In the cases reported here, a combined program of worker training on the hazards encountered during the use of 40% HF or 6% HF/15% HNO<sub>3</sub> together with the use of Hexafluorine(R) for emergent decontamination within the first 2 minutes following eye or skin exposure allowed

avoidance of the development of chemical burns and their sequelae. Hexafluorine(R) has thus been demonstrated to be efficacious for decontamination of eye or skin splashes with either concentrated or dilute HF solutions in the workplace.

Hexafluorine(R) is an HF decontamination compound which acts through a combination of physical and chemical mechanisms. It is hypertonic and can thus osmotically recover a portion of the HF which has already penetrated into the tissues. Its amphoteric properties allow it to bind both  $H^+$  and  $F^-$  ions, thus acting both against the acidity of HF and the tissue toxicity of the fluoride ion (37-39). In vitro, Hexafluorine(R) has 100 times the affinity of calcium gluconate for the fluoride ion (38).

The various treatment modalities found in the medical literature are focused on the toxic potential of HF and are based on chelation or binding of the fluoride ion, while not addressing the potential contribution of the hydrogen ion in the development of burns. Calcium gluconate has only a very weak action on the acidity of HF, making multiple topical applications or injections necessary. Treatment following splashes with hydrofluoric acid when water decontamination is followed by calcium gluconate must be repetitive, of long duration, and sometimes dependent on the re-occurrence of the sensation of pain, which may be unnecessarily uncomfortable for the patient (40-41).

By acting on all the toxic mechanisms of HF, Hexafluorine(R) allows rapid decontamination of splashes, prevention of burns and their sequelae, rapid resolution of pain, and avoidance of the need for further medical or surgical treatment. The excellent clinical outcomes in the cases reported here demonstrate both the efficacy of Hexafluorine(R) for decontamination of HF splashes and the value of a training program for workers about the hazards of the chemical and first aid procedures (42). When workers are familiar with emergency protocols for HF splash response, the duration of chemical contact can be significantly decreased.

*In vitro* studies, *in vivo* experimental animal studies, and results from 5 previously- reported human HF eye or skin splash cases, as well as the 11 cases reported here, demonstrate that Hexafluorine(R) is efficacious for HF decontamination (38-39).

### CONCLUSION

When workers have been trained about the hazards of HF splashes and rapidly utilize a decontamination compound such as Hexafluorine(R) which acts to neutralize all of the toxic mechanisms of this chemical, it is possible to prevent the development of eye or skin burns and their sequelae as demonstrated in the cases presented here. The particular value of Hexafluorine(R) for emergent decontamination of HF eye or skin splashes is that, unlike other decontamination measures, it can prevent the development of burns and the necessity for further medical or surgical treatment, potential serious or even fatal systemic toxicity, sequelae, and lost work time. It can rapidly and easily be used by the victim or co-worker directly at the accident site.

## REFERENCES

1. Dayal HH, Baranowski T, Li YH, et al: Hazardous chemicals: Psychological dimensions of the health sequelae of a community exposure in Texas. *J Epidemiol Comm Health.* 48:560-568 1994.
2. Beaudouin L, Le Trionnaire C, Nail JP: Accidents du travail dus a l'utilisation de l'acide fluorhydrique ou des fluorures alcalins en milieu acide (French). [Workplace accidents due to the use of hydrofluoric acid or alkaline fluorides in an acid mixture.] *arch. Mal. Prof.* June:403-405 1989
3. Camarasa J: A propos de 3 brûlures cutanées graves dues à l'acide fluorhydrique (French). [Regarding 3 serious cutaneous burns due to hydrofluoric acid.] *Arch Mal Prof.* April:422-425 1983
4. Tepperman PB: Fatality due to acute systemic fluoride poisoning following an hydrofluoric acid skin burn. *J Occup Med.* 22(10):691-692, 1980.
5. Sheridan RL, Ryan CM, Quinby WC, et al: Emergency management of major hydrofluoric acid exposures. *Burns.* 21:62-64, 1995.
6. Kirkpatrick JJR, Enion DS, Burd DAR: Hydrofluoric acid burns: A review. *Burns.* 21:483-493 1995
7. Chataigner D, Garnier R, Bonin C, et al: Brûlures et intoxication systémique mortelle secondaires à une projection d'acide fluorhydrique (French). [Burns and fatal systemic poisoning secondary to an hydrofluoric acid splash.] *Arch Mal Prof.* 53:3-29 1992.
8. Mullett L, Lee E, Genovese J, et al: Fatality due to fluoride poisoning following dermal contact with hydrofluoric acid in a palynology laboratory. *Ann Occup Hyg.* 40:705-710 1996.
9. Chan KM, Svancarek WP, Creer M: Fatality due to acute hydrofluoric acid exposure. *Clin. Toxicol.* 24(4):333-339 1987.

10. Noonan T, Carter EJ, Edelman PA, et al: Epidermal lipids and the natural history of hydrofluoric acid (HF) injury. *Burns*. 20:202-206 1994.
11. McCulley JP, Whitting DW, Petitt MG, et al: Hydrofluoric acid burns of the eye. *J Occup Med*. 25(6):447-450 1983.
12. Upfal M, Doyle C: Medical management of hydrofluoric acid exposure. *J. Occup Med*. 32:726-731,1990.
13. El Saadi MS, Hall AH, Hall PK, et al: Hydrofluoric acid dermal exposure. *Vet Human Toxicol*. 31(3):243-247 1989
14. Beiran I, Miller B, Bentur Y: The efficacy of calcium gluconate in ocular hydrofluoric acid burns. *Human Exp. Toxicol*. 16(4):223-228 1997.
15. Dunn BJ, MacKinnon MA, Knowlden NF, et al: Topical treatments for hydrofluoric acid dermal burns. *J Occup Environ Med*. 38:507-514 1996.
16. Boink ABTJ, Meulenbelt JAN: Systemic fluoride poisoning following dermal hydrofluoric acid exposure: Development of an intravenous sodium fluoride infusion model in rats. *J Toxicol-Cut Ocular Toxicol*. 14:75-87 1995.
17. St. Noordhoek L, Botens S, Mani MM: A study to determine the efficacy of treatments for hydrofluoric acid burns. *J Burn Care Rehabil*. 16:253-257 1995.
18. Cox RD, Osgood KA: Evaluation of intravenous magnesium sulfate for the treatment of hydrofluoric acid burns. *Clin Toxicol*. 32:23-136 1994.
19. Dowback G, Rose K, Rohrich RJ: A biochemical and histologic rationale for the treatment of hydrofluoric acid burns with calcium gluconate. *J Burn Care Rehabil*. 15:323-327 1994.
20. Burkhart KK, Brent J, Kirk MA, et al: Comparison of topical magnesium and calcium treatment for dermal hydrofluoric acid burns. *Ann Emerg Med*. 24:9-13 1994

21. Kono K, Yoshida Y, Watanabe M, et al: An experimental study on the treatment of hydrofluoric acid burns. *Arch Environ Contam Toxicol.* 22:414-418 1992.
22. Bracken WM, Cuppage F, McLaury RL, et al: Comparative effectiveness of topical treatments for hydrofluoric acid burns. *J Occup Med.* 27:733-739 1985.
23. Carney SA, Hall M, Lawrence JC, et al: Rationale of the treatment of hydrofluoric acid burns. *Br J Ind Med.* 31:317-321 1974.
24. Rubinfeld RS, Silbert DI, Arentsen JJ, et al: Ocular hydrofluoric acid burns. *Am. J. Ophthalmol.* 114:420-423 1992.
25. Matsuno K: The treatment of hydrofluoric acid burns. *Occup Med.* 46:313-317 1996.
26. Greco RJ, Hartford CE, Haith LR, et al: Hydrofluoric acid-induced hypocalcemia. *J Trauma.* 28: 1593-1596 1988.
27. Trevino MA, Herrmann GH, Sprout WL: Treatment of severe hydrofluoric acid exposures. *J Occup Med.* 25:861-863 1983.
28. Browne TD: The treatment of hydrofluoric acid burns. *J Soc Occup Med.* 24:80-89 1974.
29. Wetherhold JM, Sheperd FP: Treatment of hydrofluoric acid burns. *J Occup Med.* 7:193-195 1965.
30. Griffith FD: Hydrofluoric acid burn: Latent period was key factor. *Am Ind. Hyg Assoc J.* 48: 451-452 1987
31. Saada V, Patarin M, Sans S, et al: Nécroses cutanées à l'acide fluorhydrique (French). [Cutaneous necrosis from hydrofluoric acid.] *Ann Dermatol Venereol.* 122:512-513 1995.
32. Henry JA, Hila KK: Intravenous regional calcium gluconate perfusion for hydrofluoric acid burns. *Clin Toxicol.* 30(2):203-207 1992.
33. Chick LR, Borah G.: Calcium carbonate gel therapy for hydrofluoric acid burn of the hand. *Plastic Reconstit Surg.* 86:935-940 1990.

34. Buckingham FM: Surgery: A radical approach to severe hydrofluoric acid burns. *J Occup Med.* 30: 873-874 1988.
35. Barbier F, Bonnet P, Julie R, et al: Brûlures cutanées par acide fluorhydrique, A propos de 32 cas (French). [Hydrofluoric acid cutaneous burns. Regarding 32 cases.] *Arch. Mal. Prof.* Apr:400-402 1987
36. Saada SA, Hall M, Lawrence JC, et al: Rationale of the treatment of hydrofluoric acid burns. *Br. J. Ind Med.* 31:317-321 1974.
37. Burgher F, Blomet J, Mathieu L: *Le Risque Chimique et la Santé au Travail: Essai de Toxicologie Réflexive* (French). [Chemical Risk and Occupational Health: Review of Toxicology], pp. 610-616. Prevor, Valmondois, France 1996.
38. Josset P, Blomet J, Lym SK, et al: Theoretical and experimental evaluation of decontamination measures for burns with hydrofluoric acid. Proprietary data of Laboratoire PREVOR, Valmondois, France 1992.
39. Hall AH, Blomet J, Gross M, et al: Hexafluorine(R) for emergent decontamination of hydrofluoric acid eye/skin splashes. *Semiconductor Safety Assoc J.* 14:30-33 2000.
40. Lheureux P, Goldschmidt D, Hossey D, et al: Brûlures digitales par l'acide fluorhydrique (French). [Digital burns by hydrofluoric acid.] *Rean Soins Intens Med Urg.* 7:227-230 1991.
41. White JW: Hydrofluoric acid burns. *Cutis.* 34:241-244 1984.
42. Perrotte MC, Caubet A, Paysant F, et al: Acide fluorhydrique: Maitrise du risque à partir d'un exemple (French). [Hydrofluoric acid: An example of managing the risk.] *Arch Mal Prof.* Nov. 541-543 1993.

**Table 1.** Emergency Decontamination of 40% HF or 6% HF/15% HNO<sub>3</sub> Eye Splashes with Hexafluorine(R)

No. Cases	Exposure	Splash Area Involved	Initial Decontamination	Second Decontamination	Sequelae	Requirement for Further Treatment	Lost Work Time
1	40% HF	Eye	Hexafluorine®	Hexafluorine®	None	None	None
1	6% HF/ 15% HNO <sub>3</sub>	Eye	Hexafluorine®	Hexafluorine®	None	None	None

**Table 2.** Emergency Decontamination of 40% HF or 6% HF/15% HNO<sub>3</sub> Skin Splashes with Hexafluorine(R)

No. Cases	Exposure	Splash Area Involved (% Total body surface area)	Initial Decontamination	Second Decontamination	Sequelae	Requirement for Further Treatment	Lost Work Time
5	40% HF	0.2; 1; 4.5; 4.5; 16.5	Hexafluorine®	Hexafluorine®	None	None	None
5	6% HF/15% HNO <sub>3</sub>	0.2; 2.25; 4; 4.5; 10.5	Hexafluorine®	Hexafluorine®	None	None	None